



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON THE DIFFERENTIATION OF AN INFINITE SERIES TERM BY TERM.

BY M. B. PORTER.

IF

$$f(x) = \sum_1^{\infty} u_n(x), \quad a < x < b,$$

denote a convergent series of single valued, differentiable, real functions of the real variable x , the ordinary sufficient condition that the derivative $f'(x)$ exist and that

$$f'(x) = \sum_1^{\infty} u'_n(x),$$

namely, that each term $u_n(x)$ be differentiable and that the series $\sum_1^{\infty} u'_n(x)$ be uniformly convergent, is practically limited to the case where the uniformity can be tested by means of a so called M -series. That is, we shall have

$$f'(x) = \sum_1^{\infty} u'_n(x), \quad a < x < b,$$

if there exist a convergent series of positive terms, $\sum_1^{\infty} M_n$, such that

$$|u'_n(x)| \leq M_n, \quad a < x < b. \quad (A)$$

The practical importance of this test in that, in particular, it is immediately applicable to power series and thus leads to a simple as well as rigorous treatment of undetermined coefficients, renders an elementary proof of it desirable. The following proof, while probably not new,* has not come to the notice of the writer elsewhere, and seems to be sufficiently elementary.

If x and $x + \Delta x$ lie in the interval of convergence of the series

$$f(x) = \sum_1^{\infty} u_n(x),$$

* Dini, *Functionen einer reellen Grösse*, p. 154, uses the Mean Value Theorem, but as he deduces a more general criterion, his proof is much less elementary.

then, by the primary definition of convergence, we have

$$\frac{f(x + \Delta x) - f(x)}{\Delta x} = \sum_1^{\infty} \frac{u_n(x + \Delta x) - u_n(x)}{\Delta x},$$

and, by the first Mean Value Theorem,

$$\frac{f(x + \Delta x) - f(x)}{\Delta x} = \sum_1^{\infty} u'_n(x + \theta_n \Delta x), \quad 0 < \theta_n < 1. \quad (1)$$

If ϵ be an arbitrarily assigned positive number as small as we please, (A) tells us that a positive integer N can be found such that

$$\left| \sum_N^{\infty} u'_n(x + \theta_n \Delta x) \right| < \frac{\epsilon}{2}. \quad (2)$$

Furthermore, since each term $u_n(x)$ has a derivative $u'_n(x)$, a number δ can be found such that, if $|\Delta x| < \delta$,

$$\frac{u_n(x + \Delta x) - u_n(x)}{\Delta x} = u'_n(x + \theta_n \Delta x)$$

will differ from its limit, $u'_n(x)$, numerically by less than $\epsilon/2n$, so that

$$\left| \sum_1^n u'_n(x + \theta_n \Delta x) - \sum_1^n u'_n(x) \right| < \frac{\epsilon}{2}. \quad (3)$$

Hence it follows that

$$\left| \frac{f(x + \Delta x) - f(x)}{\Delta x} - \sum_1^{\infty} u'_n(x) \right| < \epsilon, \quad |\Delta x| < \delta,$$

which proves the theorem.

The continuity, or even the integrability, of the derivatives $u'_n(x)$ is not postulated in the above proof, since the Mean Value Theorem merely requires that the derivatives exist.

The same proof can be applied when x denotes the complex variable, $x_1 + x_2 i$, the Mean Value Theorem* for functions of a complex variable now being employed.

NEW HAVEN, CONNECTICUT,
AUGUST, 1901.

* Stolz, *Diff. u. Integralrechnung*, vol. 2, p. 95.